

## Anti-UGT1A10 antibody

<b>Cat. No.</b>	ml123478
<b>Package</b>	25 µl/100 µl/200 µl
<b>Storage</b>	-20°C, pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol

### Product overview

<b>Description</b>	Anti-UGT1A10 rabbit polyclonal antibody
<b>Applications</b>	ELISA, WB
<b>Immunogen</b>	Fusion protein of human UGT1A10
<b>Reactivity</b>	Human, Mouse
<b>Content</b>	0.5 mg/ml
<b>Host species</b>	Rabbit
<b>Ig class</b>	Immunogen-specific rabbit IgG
<b>Purification</b>	Antigen affinity purification

### Target information

<b>Symbol</b>	UGT1A10
<b>Full name</b>	UDP glucuronosyltransferase 1 family, polypeptide A10
<b>Synonyms</b>	UDPGT; UGT1J; UGT-1J; UGT1-10; UGT1.10
<b>Swissprot</b>	Q9HAW8

### Target Background

This gene encodes a UDP-glucuronosyltransferase, an enzyme of the glucuronidation pathway that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into water-soluble, excretable metabolites. This gene is part of a complex locus that encodes several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons followed by four common exons. Four of the alternate first exons are considered pseudogenes.

订购热线: 4008-898-798

**Applications**

**Western blotting**

Predicted band size:60 kDa

Positive control:231 and hela cell, mouse liver tissue

Recommended dilution: 200-1000

Gel: 6%SDS-PAGE


Lysate: 40 µg

Lane 1-3: 231 and hela cell, mouse liver tissue

Primary antibody: ml123478(UGT1A10 Antibody) at dilution 1/300

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 10 seconds



**ELISA**

Recommended dilution:1000-2000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio\_cn@yeah.net

网址: www.mlbio.cn