

Anti-RICTOR antibody

 Cat. No.
 ml121809

 Package
 25 μl/100 μl/200 μl

 Storage
 -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview	
Description	Anti-RICTOR rabbit polyclonal antibody
Applications	ELISA, IHC
Immunogen	Fusion protein of human RICTOR
Reactivity	Human, Mouse
Content	0.3 mg/ml
Host species	Rabbit
lg class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification
Target information	
Symbol	RICTOR
Full name	RPTOR independent companion of MTOR, complex 2
Synonyms	PIA; AVO3; hAVO3

Q6R327

Target Background

Swissprot

Rapamycin-insensitive companion of mTOR is a protein that in humans is encoded by the RICTOR gene. RICTOR and MTOR are components of a protein complex that integrates nutrient- and growth factor-derived signals to regulate cell growth. Subunit of mTORC2, which regulates cell growth and survival in response to hormonal signals. mTORC2 is activated by growth factors, but, in contrast to mTORC1, seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors.



订购热线: 4008-898-798

Applications Immunohistochemistry

Predicted cell location: Cytoplasm Positive control: Human thyroid cancer Recommended dilution: 25-200



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml121809(RICTOR Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: ×200)

Predicted cell location: Cytoplasm Positive control: Human liver cancer Recommended dilution: 25-200





The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ml121809(RICTOR Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: ×200)

ELISA

Recommended dilution: 1000-5000

- 联系电话: 4008-898-798, 021-61725725
- 联系QQ: 2881505695, 2881505696
- 邮箱: mlbio_cn@yeah.net 网址: www.mlbio.cn