

兔抗 CLEC7A 多克隆抗体

中文名称：兔抗 CLEC7A 多克隆抗体

英文名称：Anti-CLEC7A rabbit polyclonal antibody

别名：C-type lectin domain containing 7A; BGR; CD369; CANDF4; SCARE2; DECTIN1; CLECSF12

相关类别：一抗

储存：冷冻 (-20℃)

宿主：Rabbit

抗原：CLEC7A

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. The encoded glycoprotein is a small type II membrane receptor with an extracellular C-type lectin-like domain fold and a cytoplasmic domain with an immunoreceptor tyrosine-based activation motif. It functions as a pattern-recognition receptor that recognizes a variety of beta-1,3-linked and beta-1,6-linked glucans from fungi and plants, and in this way plays a role in innate immune response. Alternate transcriptional splice variants, encoding different isoforms, have been

	en characterized. This gene is closely linked to other CTL /CTLD superfamily members on chromosome 12p13 in the natural killer gene complex region.
Applications:	ELISA, IHC
Name of antibody:	CLEC7A
Immunogen:	Fusion protein of human CLEC7A
Full name:	C-type lectin domain containing 7A
Synonyms:	BGR; CD369; CANDF4; SCARE2; DECTIN1; CLECSF12
SwissProt:	Q9BXN2
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human tonsil
IHC Recommend dilution:	100-200

