

兔抗 AR(Phospho-Tyr363) 多克隆抗体

中文名称：兔抗 AR(Phospho-Tyr363) 多克隆抗体

英文名称： Anti-AR(Phospho-Tyr363) rabbit polyclonal antibody

别名： KD; AIS; TFM; DHTR; SBMA; HYPSP1; NR3C4; SMAX1; HUMARA

相关类别： 一抗

储存： 冷冻（-20℃） 避光

宿主： Rabbit

抗原： AR(Phospho-Tyr363)

反应种属： Human Mouse

标记物： Unconjugate

克隆类型： Unconjugate

技术规格

Background:

The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expan

	<p>sion of the polyglutamine tract causes spinal bulbar muscular atrophy (Kennedy disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Two alternatively spliced variants encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008]</p>
Applications:	WB
Name of antibody:	AR(Phospho-Tyr363)
Immunogen:	Peptide sequence around phosphorylation site of tyrosine 363 (D-Y-Y(p)-N-F) derived from Human Androgen Receptor.
Full name:	androgen receptor
Synonyms :	KD; AIS; TFM; DHTR; SBMA; HYP1; NR3C4; SMAX1; HUMARA
SwissProt:	P10275
WB Predicted band size:	99 kDa
WB Positive control:	Huvec cells, COS7 cells and JK cells lysates
WB Recommended dilution:	500-1000

