

兔抗 EIF4A2 多克隆抗体

中文名称：兔抗 EIF4A2 多克隆抗体

英文名称：Anti-EIF4A2 rabbit polyclonal antibody

别名：eukaryotic translation initiation factor 4A2; DDX2B; EIF4A; EIF4F; BM-010; eIF4A-II; eIF-4A-II

相关类别：一抗

储存：冷冻 (-20℃)

宿主：Rabbit

抗原：EIF4A2

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|--------------------------|---|
| Background: | ATP-dependent RNA helicase which is a subunit of the eIF4F complex involved in cap recognition and is required for mRNA binding to ribosome. In the current model of translation initiation, eIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which is necessary to allow efficient binding of the small ribosomal subunit, and subsequent scanning for the initiator codon. |
| Applications: | ELISA, IHC |
| Name of antibody: | EIF4A2 |

| | |
|------------------------------------|--|
| Immunogen: | Fusion protein of human EIF4A2 |
| Full name: | eukaryotic translation initiation factor 4A2 |
| Synonyms: | DDX2B; EIF4A; EIF4F; BM-010; eIF4A-II; eIF-4A-II |
| SwissProt: | Q14240 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human liver cancer and Human esophagus cancer |
| IHC Recommend dilution: | 50-200 |



