

小鼠抗 WDR77 单克隆抗体

中文名称： 小鼠抗 WDR77 单克隆抗体

英文名称： Anti-WDR77 mouse monoclonal antibody

别名： WD repeat domain 77; p44; MEP50; MEP-50; HKMT1069; Nbla10071; p44/Mep50

相关类别： 一抗

储存： 冷冻（-20℃）

宿主： Mouse

抗原： WDR77

反应种属： Human, Mouse, Rat


标记物： Unconjugate

克隆类型： Mouse Monoclonal

技术规格

Background:	The protein encoded by this gene is an androgen receptor coactivator that forms a complex with protein arginine methyltransferase 5, which modifies specific arginines to dimethylarginines in several spliceosomal Sm proteins. The encoded protein may be involved in the early stages of prostate cancer, with most of the protein being nuclear-localized in benign cells but cytoplasmic in cancer cells. Several transcript variants encoding different isoforms have been found for this gene.
Applications:	WB, IF
Name of antibody:	WDR77
Immunogen:	Fusion protein of human WDR77
Full name:	WD repeat domain 77

Synonyms:	p44; MEP50; MEP-50; HKMT1069; Nbla10071; p44/Mep50
SwissProt:	Q9BQA1
WB Predicted band size:	37 kDa
WB Positive control:	C6, K562 and 3T3 cell lysates
WB Recommended dilution:	1000-5000
IF positive control:	Hela cells
IF Recommend dilution:	50-100



mlbio 梅联生物
Good elisakit producers

