

订购热线: 4008-898-798

Anti-DDR2 antibody

Cat. No. ml160967

Package 25 μl/100 μl/200 μl

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-DDR2 rabbit polyclonal antibody

Applications ELISA, IHC

Immunogen Synthetic peptide of human DDR2

Reactivity Human, Mouse
Content 0.3 mg/ml
Host species Rabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol DDR2

Full name discoidin domain receptor tyrosine kinase 2

Synonyms TKT; MIG20a; NTRKR3; TYRO10

Swissprot Q16832

Target Background

Receptor tyrosine kinases (RTKs) play a key role in the communication of cells with their microenvironment. These molecules are involved in the regulation of cell growth, differentiation, and metabolism. In several cases the biochemical mechanism by which RTKs transduce signals across the membrane has been shown to be ligand induced receptor oligomerization and subsequent intracellular phosphorylation. This autophosphorylation leads to phosphorylation of cytosolic targets as well as association with other molecules, which are involved in pleiotropic effects of signal transduction. RTKs have a tripartite structure with extracellular, transmembrane, and cytoplasmic regions. This gene encodes a member of a novel subclass of RTKs and contains a distinct extracellular region encompassing a factor VIII-like domain.





订购热线: 4008-898-798


Applications

Immunohistochemistry

Predicted cell location: Cytoplasm Positive control: Human thyroid cancer







The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml160967(DDR2 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: ×200)

ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net 网址: www.mlbio.cn