

Anti-HCAR2 antibody

Cat. No.	ml161722
Package	25 µl/100 µl/200 µl
Storage	-20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol

Product overview

Description	Anti-HCAR2 rabbit polyclonal antibody
Applications	ELISA, IHC
Immunogen	Synthetic peptide of human HCAR2
Reactivity	Human, Mouse, Rat
Content	0.8 mg/ml
Host species	Rabbit
Ig class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification

Target information

Symbol	HCAR2
Full name	hydroxycarboxylic acid receptor 2
Synonyms	HCA2; HM74a; HM74b; PUMAG; NIACR1; Puma-g; GPR109A
Swissprot	Q8TDS4

Target Background

HCAR2, also known as PUMAG or Puma-g, is a member of the G protein coupled receptor (GPCR) superfamily. HCAR2 is highly expressed in adipocytes, immune cells and spleen. Like all members of the GPCR superfamily, HCAR2 contains seven transmembrane domains.

订购热线: 4008-898-798



Applications

Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human gasrtic cancer

Recommended dilution: 100-300





The image on the left is immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using ml161722(HCAR2 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: ×200)

Predicted cell location: Cytoplasm

Positive control: Human thyroid cancer

Recommended dilution: 100-300



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml161722(HCAR2 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: ×200)

ELISA

Recommended dilution: 2000-10000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn