

订购热线: 4008-898-798

Anti-CORO2A antibody

Cat. No. ml163577

Package 25 μl/100 μl/200 μl

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-CORO2A rabbit polyclonal antibody

Applications ELISA, IHC

Immunogen Synthetic peptide of human CORO2A

ReactivityHumanContent1.8 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol CORO2A
Full name coronin 2A

Synonyms IR10; WDR2; CLIPINB

Swissprot Q92828

Target Background

This gene encodes a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. This protein contains 5 WD repeats, and has a structural similarity with actin-binding proteins: the D. discoideum coronin and the human p57 protein, suggesting that this protein may also be an actin-binding protein that regulates cell motility. Alternative splicing of this gene generates 2 transcript variants.




订购热线: 4008-898-798

Applications

Immunohistochemistry

Predicted cell location: Cytoplasm and Cell membrane

Positive control: Human gastric cancer Recommended dilution: 30-150





The image on the left is immunohistochemistry of paraffin-embedded Human gastric cancer tissue using ml163577(CORO2A Antibody) at dilution 1/45, on the right is treated with synthetic peptide. (Original magnification: ×200)

ELISA

Recommended dilution: 5000-10000


联系电话: 4008-898-798, 021-61725725


联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net 网址: www.mlbio.cn

Predicted cell location: Cytoplasm and Cell membrane

Positive control: Human liver cancer Recommended dilution: 30-150





The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ml163577(CORO2A Antibody) at dilution 1/45, on the right is treated with synthetic peptide. (Original magnification: ×200)