

订购热线: 4008-898-798

Anti-MBOAT4 antibody

Cat. No. ml163989

Package 25 μ l/100 μ l/200 μ l

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-MBOAT4 rabbit polyclonal antibody

Applications ELISA, IHC

Immunogen Synthetic peptide of human MBOAT4

ReactivityHuman, MouseContent0.9 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol MBOAT4

Full name membrane bound O-acyltransferase domain containing 4

Synonyms GOAT; OACT4; FKSG89

Swissprot Q96T53

Target Background

Mediates the octanoylation of ghrelin at 'Ser-3'. Can use a variety of fatty acids as substrates including octanoic acid, decanoic acid and tetradecanoic acid.



订购热线: 4008-898-798


Applications


Immunohistochemistry

Predicted cell location: Endoplasmic reticulum membrane

Positive control: Human cervical cancer

Recommended dilution: 30-150





The image on the left is immunohistochemistry of paraffin-embedded Human cervical cancer tissue using ml163989(MBOAT4 Antibody) at dilution 1/30, on the right is treated with synthetic peptide. (Original magnification: ×200)

ELISA

Recommended dilution: 5000-10000

联系电话: 4008-898-798, 021-61725725


联系QQ: 2881505695, 2881505696


邮箱: mlbio_cn@yeah.net 网址: www.mlbio.cn

Predicted cell location: Endoplasmic reticulum membrane

Positive control: Human colorectal cancer

Recommended dilution: 30-150





The image on the left is immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using ml163989(MBOAT4 Antibody) at dilution 1/30, on the right is treated with synthetic peptide. (Original magnification: ×200)