

## Anti-CDKN2AIP antibody

<b>Cat. No.</b>	ml221080
<b>Package</b>	25 µl/100 µl/200 µl
<b>Storage</b>	-20°C, pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol

### Product overview

<b>Description</b>	Anti-CDKN2AIP rabbit polyclonal antibody
<b>Applications</b>	ELISA, IHC
<b>Immunogen</b>	Fusion protein of human CDKN2AIP
<b>Reactivity</b>	Human
<b>Content</b>	0.2 mg/ml
<b>Host species</b>	Rabbit
<b>Ig class</b>	Immunogen-specific rabbit IgG
<b>Purification</b>	Antigen affinity purification

### Target information

<b>Symbol</b>	CDKN2AIP
<b>Full name</b>	CDKN2A interacting protein
<b>Synonyms</b>	CARF
<b>Swissprot</b>	Q9NXV6

### Target Background

CDKN2AIP (CDKN2A-interacting protein), also known as CARF, is a 580 amino acid protein that activates p53 via p14 ARF (alternate reading frame)-dependent and independent pathways. CDKN2AIP-dependent activation of p53, a protein that up-regulates growth arrest and apoptosis-related genes in response to stress signals, leads to an enhancement of p53 function. Expression levels of CDKN2AIP and p53 show an inverse relationship that is caused by a negative-feedback control via a proteasome-mediated degradation pathway. CDKN2AIP is expressed ubiquitously across tissue samples and, along with p14 ARF, is localized to the perinucleolar region within the nucleus.

订购热线: 4008-898-798


#### Applications

##### Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human brain

Recommended dilution: 25-100



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using ml221080(CDKN2AIP Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification:  $\times 200$ )

##### ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: [mlbio\\_cn@yeah.net](mailto:mlbio_cn@yeah.net)

网址: [www.mlbio.cn](http://www.mlbio.cn)