

兔抗 CIDEC 多克隆抗体

中文名称: 兔抗 CIDEC 多克隆抗体

英文名称: Anti-CIDEC rabbit polyclonal antibody

别 名: CIDE3; FPLD5; FSP27; CIDE-3

相关类别: 一抗

抗 原: CIDEC

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human, Mouse, Rat

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

	This gene encodes a member of the cell death-inducing DNA fragmentation factor-like effector family. Members of this family play important roles in apoptosis. The encoded protein promotes lipid droplet formation in adipocytes and may mediate adipocyte apoptosis. This gene is
Background:	s regulated by insulin and its expression is positively co
	rrelated with insulin sensitivity. Mutations in this gene
	may contribute to insulin resistant diabetes. A pseudoge
	ne of this gene is located on the short arm of chromos
	ome 3. Alternatively spliced transcript variants that enco
	de different isoforms have been observed for this gene.
Applications:	ELISA, WB, IHC



Name of antibody:	CIDEC
Immunogen:	Fusion protein of human CIDEC
Full name:	cell death inducing DFFA like effector c
Synonyms:	CIDE3; FPLD5; FSP27; CIDE-3
SwissProt:	Q96AQ7
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human esophagus cancer
IHC Recommend dilution:	50-100
WB Predicted band size:	27 kDa
WB Positive control:	Mouse small intestines tissue lysate
WB Recommended dilution:	500-2000





