

## CLASP2 抗原(重组蛋白)

- 中文名称: CLASP2 抗原(重组蛋白)
- 英文名称: CLASP2 Antigen (Recombinant Protein)
- 别 名: cytoplasmic linker associated protein 2
- 储存: 冷冻 (-20℃)
- 相关类别: 抗原

概述

Fusion protein corresponding to a region derived from 232-431 amino acids of human CLASP2

## 技术规格

| Full name:         | cytoplasmic linker associated protein 2  |
|--------------------|--|
| Swissprot:         | 075122   |
| Gene Accession:    | BC029035   |
| Purity:            | >85%, as determined by Coomassie blue stained SDS-PAGE   |
| Expression system: | Escherichia coli   |
| Tags:              | His tag C-Terminus, GST tag N-Terminus   |
| Background:        | CLASP2 (cytoplasmic linker associated protein 2), also known as KIAA 0627 or hOrbit2, is a 1,294 amino acid protein that contains five HEA T repeats and localizes to the cytoplasm and the cytoskeleton, as wel I as to the kinetochore and the golgi apparatus. Expressed primarily i n brain tissue, CLASP2 functions as a microtubule plus-end tracking p rotein that regulates the stability of dynamic microtubules and is req uired for the proper polarization of cytoplasmic microtubule arrays in migrating cells. CLASP2 interacts with EB1, EB3, ELKS and CLIP-115 an d, in addition to stabilizing microtubules, plays an important role in |



| maintaining the stability of the kinetochore and is crucial for proper          |
|---|
| chromosomal alignment. CLASP2 is subject to phosphorylation by GS               |
| K-3 $\beta$ , an event that is thought to negatively regulate the ability of CL |
| ASP2 to bind to microtubules. Two isoforms of CLASP2, designated b              |
| eta and gamma, exist due to alternative splicing events.                        |