

兔抗 ASCC2 多克隆抗体

- 中文名称：兔抗 ASCC2 多克隆抗体
- 英文名称：Anti-ASCC2 rabbit polyclonal antibody
- 别名：p100; ASC1p100
- 相关类别：一抗
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 抗原：ASCC2
- 反应种属：Human, Mouse
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:

ASCC2 (activating signal cointegrator 1 complex subunit 2), also known as ASC1P100, is a 757 amino acid protein that is ubiquitously expressed and contains one CUE domain. The gene encoding ASCC2 maps to human chromosome 22, which houses over 500 genes and is the second smallest human chromosome. Mutations in several of the genes that map to chromosome 22 are involved in the development of Phelan-McDermid syndrome, Neurofibromatosis type 2, autism and schizophrenia. Additionally, translocations between chromosomes 9 and 22 may lead to the formation of the Philadelphia Chrom

	osome and the subsequent production of the novel fusion protein BCR-Abl, a potent cell proliferation activator found in several types of leukemias.
Applications:	ELISA, WB
Name of antibody:	ASCC2
Immunogen:	Synthetic peptide of human ASCC2
Full name:	activating signal cointegrator 1 complex subunit 2
Synonyms :	p100; ASC1p100
SwissProt:	Q9H1I8
ELISA Recommended dilution:	2000-5000
WB Predicted band size:	86 kDa
WB Positive control:	A431, 293T and hela cells
WB Recommended dilution:	500-2000

