

CST1 抗原(重组蛋白)

中文名称: CST1 抗原(重组蛋白)

英文名称: CST1 Antigen (Recombinant Protein)

别 名: cystatin SN

储 存: 冷冻(-20℃)

相关类别: 抗原

概述

Fusion protein corresponding to a region derived from 21-141 amino acids of human CST1

技术规格

Full name:	cystatin SN
Swissprot:	P01037
Gene Accession:	BC021225
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine pro tease inhibitors, while others have lost or perhaps never acquired thi s inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes



and pseudogenes. This gene is located in the cystatin locus and enc odes a cysteine proteinase inhibitor found in saliva, tears, urine, and seminal fluid.