

兔抗 BAG2 多克隆抗体

中文名称：兔抗 BAG2 多克隆抗体

英文名称： Anti-BAG2 rabbit polyclonal antibody

别名： BAG-2; dJ41711.2

相关类别： 一抗

储存： 冷冻（-20℃）

宿主： Rabbit

抗原： BAG2

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:	BAG proteins compete with Hip for binding to the Hsc70 /Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The predicted BAG2 protein contains 211 amino acids. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner.
Applications:	ELISA, WB, IHC

Name of antibody:	BAG2
Immunogen:	Synthetic peptide of human BAG2
Full name:	BCL2-associated athanogene 2
Synonyms :	BAG-2; dJ41711.2
SwissProt:	O95816
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human tonsil and human esophagus cancer
IHC Recommend dilution:	30-150
WB Predicted band size:	24 kDa
WB Positive control:	Human fetal muscle tissue , HEPG2 , Jurkat,Hela , A431 and A549 cell
WB Recommended dilution:	500-2000



