

兔抗 BAG3 多克隆抗体

中文名称：兔抗 BAG3 多克隆抗体

英文名称： Anti-BAG3 rabbit polyclonal antibody

别名： BIS; BAG-3; CAIR-1

抗原： BAG3

储存： 冷冻（-20℃）

宿主： Rabbit

相关类别： 一抗

反应种属： Human, Mouse

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The protein encoded by this gene contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone

	ne activity in a Hip-repressible manner.
Applications:	ELISA, WB, IHC
Name of antibody:	BAG3
Immunogen:	Fusion protein of human BAG3
Full name:	BCL2 associated athanogene 3
Synonyms:	BIS; MFM6; BAG-3; CAIR-1
SwissProt:	O95817
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human cervical cancer and Human colorectal cancer
IHC Recommend dilution:	40-200
WB Predicted band size:	62 kDa
WB Positive control:	Mouse heart tissue lysate
WB Recommended dilution:	500-2000



