

## 兔抗 CTLA4 多克隆抗体

中文名称：兔抗 CTLA4 多克隆抗体

英文名称：Anti-CTLA4 rabbit polyclonal antibody

别名：cytotoxic T-lymphocyte associated protein 4; CD; GSE; GRD4; ALPS5; CD152; CTLA-4; IDDM12; CELIAC3

相关类别：一抗

抗原：CTLA4

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

**Background:**

This gene is a member of the immunoglobulin superfamily and encodes a protein which transmits an inhibitory signal to T cells. The protein contains a V domain, a transmembrane domain, and a cytoplasmic tail. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. The membrane-bound isoform functions as a homodimer interconnected by a disulfide bond, while the soluble isoform functions as a monomer. Mutations in this gene have been associated with insulin-dependent diabetes mellitus, Graves disease, Hash

|                                    |   |
|------------------------------------|---|
|                                    | imoto thyroiditis, celiac disease, systemic lupus erythematosus, thyroid-associated orbitopathy, and other autoimmune diseases. |
| <b>Applications:</b>               | ELISA, IHC  |
| <b>Name of antibody:</b>           | CTLA4   |
| <b>Immunogen:</b>                  | Synthetic peptide of human CTLA4  |
| <b>Full name:</b>                  | cytotoxic T-lymphocyte associated protein 4   |
| <b>Synonyms:</b>                   | CD; GSE; GRD4; ALPS5; CD152; CTLA-4; IDDM12; CELIAC 3   |
| <b>SwissProt:</b>                  | P16410  |
| <b>ELISA Recommended dilution:</b> | 5000-10000  |
| <b>IHC positive control:</b>       | Human liver cancer and Human ovarian cancer   |
| <b>IHC Recommend dilution:</b>     | 25-100  |



