

## 兔抗 BCR (Phospho-Tyr177)多克隆抗体

中文名称：兔抗 BCR (Phospho-Tyr177)多克隆抗体

英文名称：Anti-BCR (Phospho-Tyr177) rabbit polyclonal antibody

别名：ALL; CML; PHL; BCR1; D22S11; D22S662

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：BCR (Phospho-Tyr177)

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

**Background:**

A reciprocal translocation between chromosomes 2 and 9 produces the Philadelphia chromosome, which is often found in patients with chronic myelogenous leukemia. The chromosome 22 breakpoint for this translocation is located within the BCR gene. The translocation produces a fusion protein which is encoded by sequence from both BCR and ABL, the gene at the chromosome 9 breakpoint. Although the BCR-ABL fusion protein has been extensively studied, the function of the normal BCR gene product is not clear. The protein has serine/threonine ki

	nase activity and is a GTPase-activating protein for p21rac. Two transcript variants encoding different isoforms have been found for this gene.
<b>Applications:</b>	WB, IHC
<b>Name of antibody:</b>	BCR (Phospho-Tyr177)
<b>Immunogen:</b>	Synthetic peptide of human BCR (Phospho-Tyr177)
<b>Full name:</b>	breakpoint cluster region (Phospho-Tyr177)
<b>Synonyms :</b>	ALL; CML; PHL; BCR1; D22S11; D22S662
<b>SwissProt:</b>	P11274
<b>IHC positive control:</b>	Human tonsil tumor
<b>IHC Recommend dilution:</b>	50-100
<b>WB Predicted band size:</b>	210 kDa
<b>WB Positive control:</b>	K562 cells untreated or treated with H2O2
<b>WB Recommended dilution:</b>	500-1000



