

兔抗 BCR 多克隆抗体

中文名称：兔抗 BCR 多克隆抗体

英文名称：Anti-BCR rabbit polyclonal antibody

别名：ALL; CML; PHL; BCR1; D22S11; D22S662

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：BCR

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

A reciprocal translocation between chromosomes 22 and 9 produces the Philadelphia chromosome, which is often found in patients with chronic myelogenous leukemia. The chromosome 22 breakpoint for this translocation is located within the BCR gene. The translocation produces a fusion protein which is encoded by sequence from both BCR and ABL, the gene at the chromosome 9 breakpoint. Although the BCR-ABL fusion protein has been extensively studied, the function of the normal BCR gene product is not clear. The protein has serine/threonine kinase activity and is a GTPase-act

	tivating protein for p21rac. Two transcript variants encoding different isoforms have been found for this gene.
Applications:	ELISA, IHC
Name of antibody:	BCR
Immunogen:	Synthetic peptide of human BCR
Full name:	breakpoint cluster region
Synonyms :	ALL; CML; PHL; BCR1; D22S11; D22S662
SwissProt:	P11274
ELISA Recommended dilution:	1000-2000
IHC positive control:	Human thyroid cancer and Human liver cancer
IHC Recommend dilution:	25-100

