

## 兔抗 CYP2A13 多克隆抗体

中文名称：兔抗 CYP2A13 多克隆抗体

英文名称：Anti-CYP2A13 rabbit polyclonal antibody

别名：cytochrome P450 family 2 subfamily A member 13 CPAD; CYP2A; CYP11A13

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：CYP2A13

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

**Background:**

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum. Although its endogenous substrate has not been determined, it is known to metabolize 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol, a major nitrosamine specific to tobacco. This gene is part of a large cluster of cytochrome P450 genes from the CYP2A, CYP2B and CYP2F subfamilies on chrom

|                                    |  |
|------------------------------------|--|
|                                    | osome 19q.                                     |
| <b>Applications:</b>               | ELISA, WB                                      |
| <b>Name of antibody:</b>           | CYP2A13  |
| <b>Immunogen:</b>                  | Synthetic peptide of human CYP2A13             |
| <b>Full name:</b>                  | cytochrome P450 family 2 subfamily A member 13 |
| <b>Synonyms:</b>                   | CPAD; CYP2A; CYP11A13                          |
| <b>SwissProt:</b>                  | Q16696   |
| <b>ELISA Recommended dilution:</b> | 5000-10000                                     |
| <b>WB Predicted band size:</b>     | 57 kDa   |
| <b>WB Positive control:</b>        | A549 cell lysates                              |
| <b>WB Recommended dilution:</b>    | 200-1000                                       |



