

兔抗 BMF 多克隆抗体

中文名称：兔抗 BMF 多克隆抗体

英文名称： Anti-BMF rabbit polyclonal antibody

相关类别： 一抗

储 存： 冷冻（-20℃）

宿 主： Rabbit

抗 原： BMF

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

| | |
|----------------------|---|
| Background: | The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein contains a single BCL2 homology domain 3 (BH3), and has been shown to bind BCL2 proteins and function as an apoptotic activator. This protein is found to be sequestered to myosin V motors by its association with dynein light chain 2, which may be important for sensing intracellular damage and triggering apoptosis. Alternatively spliced transcript variants encoding different isoforms have been identified. |
| Applications: | ELISA, WB, IHC |

| | |
|------------------------------------|---|
| Name of antibody: | BMF |
| Immunogen: | Synthetic peptide of human BMF |
| Full name: | Bcl2 modifying factor |
| SwissProt: | Q96LC9 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human liver cancer and Human gastric cancer |
| IHC Recommend dilution: | 50-200 |
| WB Predicted band size: | 20 kDa |
| WB Positive control: | 293T cell lysate |
| WB Recommended dilution: | 500-2000 |



