

## 兔抗 C1QTNF3 多克隆抗体

- 中文名称：兔抗 C1QTNF3 多克隆抗体
- 英文名称：Anti-C1QTNF3 rabbit polyclonal antibody
- 别名：CORS; CORCS; CTRP3; CORS26; C1ATNF3; CORS-26
- 相关类别：一抗
- 抗原：C1QTNF3
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 反应种属：Human, Mouse
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

### 技术规格

<b>Background:</b>	C1qTNF3 (Complement C1q TNF-related protein 3/CTRP3; also CORS26 and cartanectin) is a 30-32 kDa, secreted member of the C1q and TNF-related protein (CTRP) superfamily of molecules. It is expressed by a wide variety of cells, including smooth muscle cells, fibroblasts, adipocytes, monocytes and proliferating chondrocytes. C1qTNF3 is an anti-inflammatory agent that apparently blocks LPS activation of mononuclear cells. It also has marked proliferative activity on diverse cell types such as vascular smooth muscle, chondrocytes, and endothelium. Finally, C1qTNF3 is known to act on hepatocytes and suppress hep
--------------------	---

	<p>atocyte gluconeogenesis. Mature human C1qTNF3 is 224 amino acids (aa) in length (aa 23-246). It possesses an N-terminal collagen-like domain (aa 51-113) followed by a C-terminal globular region (aa 113-246). C1qTNF3 is monomeric when intracellular, but forms a 90 kDa homotrimer plus higher-order oligomer when secreted. There are at least two potential isoform variants. One is 40-42 kDa, glycosylated, and contains a 73 aa insertion after Glu28, while a second shows concurrent deletions of aa 46-69 and 82-105. The longer 40 kDa isoform is reported to form heterotrimers and oligomers with the standard 30 kDa isoform. This has the effect of protecting the standard isoform from proteolysis. Over aa 24-246, human C1qTNF3 shares 99% aa sequence identity with mouse C1qTNF3.</p>
<b>Applications:</b>	ELISA, WB, IHC
<b>Name of antibody:</b>	C1QTNF3
<b>Immunogen:</b>	Fusion protein of human C1QTNF3
<b>Full name:</b>	C1q and tumor necrosis factor related protein 3
<b>Synonyms :</b>	CORS; CORCS; CTRP3; CORS26; C1ATNF3; CORS-26
<b>SwissProt:</b>	Q9BXJ4
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human liver cancer and human esophagus cancer
<b>IHC Recommend dilution:</b>	25-100
<b>WB Predicted band size:</b>	27 kDa
<b>WB Positive control:</b>	Human colorectal cancer tissue
<b>WB Recommended dilution:</b>	200-1000



