

兔抗 DLEC1 多克隆抗体

中文名称: 兔抗 DLEC1 多克隆抗体

英文名称: Anti-DLEC1 rabbit polyclonal antibody

别 名: CLL1; MICL; CD371; CLL-1; DCAL-2

相关类别: 一抗

抗 原: DLEC1

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

The cytogenetic location of this gene is 3p21.3, and it is located in a region that is commonly deleted in a variety of malignancies. Down-regulation of this gene has been observed in several human cancers including lung, esophageal, renal tumors, and head and neck squamous cell carcinoma. In some cases, reduced expression of this gene in tumor cells is a result of aberrant promoter methylation. Several alt ernatively spliced transcripts have been observed that contain disrupted coding regions and likely encode nonfunctional proteins.



Applications:	ELISA, IHC
Name of antibody:	DLEC1
Immunogen:	Synthetic peptide corresponding to a region derived
	from 1727-1743 amino acids of human DLEC1
Full name:	deleted in lung and esophageal cancer 1
Synonyms:	F56; DLC1; DLC-1; CFAP81
SwissProt:	Q9Y238
ELISA Recommended dilution:	5000-10000
IHC Positive control:	Human esophagus cancer;Human colorectal cancer
IHC Recommended dilution:	10-50





