

兔抗 CCL8 多克隆抗体

- 中文名称: 兔抗 CCL8 多克隆抗体
- 英文名称: Anti-CCL8 rabbit polyclonal antibody
- 别 名: C-C motif chemokine ligand 8; HC14; MCP2; MCP-2; SCYA8; SCYA10
- 相关类别: 一抗
- 储 存: 冷冻 (-20℃)
- 抗 原: CCL8
- 宿 主: Rabbit
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	This antimicrobial gene is one of several chemokine gene s clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immu noregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. Thi s chemokine is a member of the CC subfamily which is c haracterized by two adjacent cysteine residues. This cytoki ne displays chemotactic activity for monocytes, lymphocyt
	ne displays chemotactic activity for monocytes, lymphocyt es, basophils and eosinophils. By recruiting leukocytes to sites of inflammation this cytokine may contribute to tum



	or-associated leukocyte infiltration and to the antiviral sta te against HIV infection.
Applications:	elisa, ihc
Name of antibody:	CCL8
Immunogen:	Synthetic peptide of human CCL8
Full name:	C-C motif chemokine ligand 8
Synonyms:	HC14; MCP2; MCP-2; SCYA8; SCYA10
SwissProt:	P80075
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human cervical cancer
IHC Recommend dilution:	25-100

