

兔抗 CCL13 多克隆抗体

中文名称：兔抗 CCL13 多克隆抗体

英文名称： Anti-CCL13 rabbit polyclonal antibody

别名： NCC1, CKb10, MCP-4, NCC-1, SCYL1, SCYA13

抗原： CCL13

储存： 冷冻（-20℃）

宿主： Rabbit

相关类别： 一抗

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

This antimicrobial gene is one of several Cys-Cys (C C) cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils, but not neutrophils. This chemokine plays a role in accumulation of leukocytes during inflammation. It may also be involved in the recruitment of monocytes into th

	e arterial wall during atherosclerosis.
Applications:	ELISA, WB, IHC
Name of antibody:	CCL13
Immunogen:	Fusion protein of human CCL13
Full name:	C-C motif chemokine ligand 13
Synonyms:	NCC1; CKb10; MCP-4; NCC-1; SCYL1; SCYA13
SwissProt:	Q99616
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	80-400
WB Predicted band size:	11 kDa
WB Positive control:	Human ileum tissue lysate
WB Recommended dilution:	500-2000



