

## 兔抗 CCL14 多克隆抗体

中文名称: 兔抗 CCL14 多克隆抗体

英文名称: Anti-CCL14 rabbit polyclonal antibody

别名: CC-1; CC-3; CKB1; MCIF; NCC2; SY14; HCC-1; HCC-3; NCC-2; SCYL2; SCYA14; HCC-1(1-74); HCC-1/HCC-3

储存: 冷冻 (-20℃)

抗原: CCL14

宿主: Rabbit

相关类别: 一抗

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

### 技术规格

**Background:**

This gene, chemokine (C-C motif) ligand 14, is one of several CC cytokine genes clustered on 17q11.2. The CC cytokines are secreted proteins characterized by two adjacent cysteines. The cytokine encoded by this gene induces changes in intracellular calcium concentration and enzyme release in monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Read-through transcripts are also expressed that include exons from the upstream cytokine gene, chemokine (C-C mo

	tif) ligand 15, and are represented as GeneID: 348249.
<b>Applications:</b>	ELISA, IHC
<b>Name of antibody:</b>	CCL14
<b>Immunogen:</b>	Synthetic peptide of human CCL14
<b>Full name:</b>	chemokine (C-C motif) ligand 14
<b>Synonyms:</b>	CC-1; CC-3; CKB1; MCIF; NCC2; SY14; HCC-1; HCC-3 ; NCC-2; SCYL2; SCYA14; HCC-1(1-74); HCC-1/HCC-3
<b>SwissProt:</b>	Q16627
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human cervical cancer
<b>IHC Recommend dilution:</b>	20-100

