

兔抗 CCL15 多克隆抗体

中文名称：兔抗 CCL15 多克隆抗体

英文名称： Anti-CCL15 rabbit polyclonal antibody

别名： LKN1, NCC3, SY15, HCC-2, LKN-1, MIP-5, NCC-3, SCYL3, MIP-1D, MRP-2B, SCYA15, HMRP-2B

相关类别： 一抗

抗原： CCL15

储存： 冷冻（-20℃）

宿主： Rabbit

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

This gene, chemokine (C-C motif) ligand 15, is one of several CC cytokine genes clustered on 17q11.2. The CC cytokines are secreted proteins characterized by two adjacent cysteines. The cytokine encoded by this gene is chemotactic for T cells and monocytes and induces N-acetyl-beta-D-glucosaminidase release in monocytes. It induces changes in intracellular calcium concentration in monocytes and is thought to act through the CCR1 receptor. Read-through transcripts are expressed that include exons from th

	e downstream cytokine gene, chemokine (C-C motif) ligand 14, and are represented as GeneID: 348249.
Applications:	ELISA, IHC
Name of antibody:	CCL15
Immunogen:	Synthetic peptide of human CCL15
Full name:	Chemokine (C-C motif) ligand 15
Synonyms :	LKN1, NCC3, SY15, HCC-2, LKN-1, MIP-5, NCC-3, SCYL3, MIP-1D, MRP-2B, SCYA15, HMRP-2B
SwissProt:	Q16663
ELISA Recommended dilution:	1000-5000
IHC positive control:	Human breast cancer
IHC Recommend dilution:	25-100

