

兔抗 CCL16 多克隆抗体

中文名称：兔抗 CCL16 多克隆抗体

英文名称：Anti-CCL16 rabbit polyclonal antibody

别名：C-C motif chemokine ligand 16 LEC; LMC; NCC4; CKb12; HCC-4; LCC-1; Mtn-1; NCC-4; SCYL4; ILINCK; SCYA16

相关类别：一抗

抗原：CCL16

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for lymphocytes and monocytes but not for neutrophils. This cytokine also shows a potent myelosuppressive activity and suppresses proliferation of myeloid progenitor cells. The expression of this gene is upregulated by IL-10.

Applications:	ELISA, IHC
Name of antibody:	CCL16
Immunogen:	Synthetic peptide of human CCL16
Full name:	C-C motif chemokine ligand 16
Synonyms:	LEC; LMC; NCC4; CKb12; HCC-4; LCC-1; Mtn-1; NCC-4; S CYL4; ILINCK; SCYA16
SwissProt:	O15467
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer;
IHC Recommend dilution:	30-150

