

兔抗 CCL22 多克隆抗体

中文名称：兔抗 CCL22 多克隆抗体

英文名称： Anti-CCL22 rabbit polyclonal antibody

别名： C-C motif chemokine ligand 22; MDC; ABCD-1; SCYA22; STCP-1; DC/B-CK; A-152E5.1

相关类别： 一抗

储存： 冷冻（-20℃）

宿主： Rabbit

抗原： CCL22

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, dendritic cells, natural killer cells and for chronically activated T lymphocytes. It also displays a mild activity for primary activated T lymphocytes and has no chemoattractant activity for neutrophils, eosinophils and resting T lymphocytes.

	mphocytes. The product of this gene binds to chemokine receptor CCR4. This chemokine may play a role in the trafficking of activated T lymphocytes to inflammatory sites and other aspects of activated T lymphocyte physiology.
Applications:	ELISA, IHC
Name of antibody:	CCL22
Immunogen:	Synthetic peptide of human CCL22
Full name:	C-C motif chemokine ligand 22
Synonyms:	MDC; ABCD-1; SCYA22; STCP-1; DC/B-CK; A-152E5.1
SwissProt:	O00626
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human ovarian cancer
IHC Recommend dilution:	25-50

