

兔抗 CCL27 多克隆抗体

中文名称：兔抗 CCL27 多克隆抗体

英文名称：Anti-CCL27 rabbit polyclonal antibody

别名：C-C motif chemokine ligand 27; ALP; ILC; CTAK; CTACK; PESKY; ESKINE; SCYA27

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：CCL27

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene is one of several CC cytokine genes clustered on the p-arm of chromosome 9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The protein encoded by this gene is chemotactic for skin-associated memory T lymphocytes. This cytokine may also play a role in mediating homing of lymphocytes to cutaneous sites. It specifically binds to chemokine receptor 10 (CCR10). Studies of a similar murine protein indicate that these protein-receptor interactions have

	e a pivotal role in T cell-mediated skin inflammation.
Applications:	ELISA, IHC
Name of antibody:	CCL27
Immunogen:	Synthetic peptide of human CCL27
Full name:	C-C motif chemokine ligand 27
Synonyms:	ALP; ILC; CTAK; CTACK; PESKY; ESKINE; SCYA27
SwissProt:	Q9Y4X3
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	25-100

