

兔抗 CCNC(Ab-275) 多克隆抗体

中文名称：兔抗 CCNC(Ab-275) 多克隆抗体

英文名称：Anti-CCNC(Ab-275) rabbit polyclonal antibody

别名：CycC

相关类别：一抗

储存：冷冻（-20℃）避光

抗原：CCNC(Ab-275)

宿主：Rabbit

反应种属：Human Mouse

标记物：Unconjugate

克隆类型：Unconjugate

技术规格

Background:

Component of the Mediator complex, a coactivator involved in regulated gene transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. Binds to and activates cyclin-dependent kinase CDK 8 that phosphorylates the CTD (C-terminal domain) of th

	e large subunit of RNA polymerase II (RNAP II), which may inhibit the formation of a transcription initiation complex.
Applications:	WB
Name of antibody:	CCNC(Ab-275)
Immunogen:	Synthesized non-phosphopeptide derived from human Cyclin C around the phosphorylation site of serine 275 (N-G-S(p)-Q-N).
Full name:	cyclin C
Synonyms :	CycC
SwissProt:	P24863
WB Predicted band size:	33 kDa
WB Positive control:	LOVO cells lysate
WB Recommended dilution:	500-3000

