

兔抗 EXOG 多克隆抗体

中文名称：兔抗 EXOG 多克隆抗体

英文名称：Anti-EXOG rabbit polyclonal antibody

别名：exo/endonuclease G; ENGL; ENGLA; ENGLB; ENGL-a; ENGL-b; ENDOGL1; ENDOGL2

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：EXOG

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes an endo/exonuclease with 5'-3' exonuclease activity. The encoded enzyme catalyzes the hydrolysis of ester linkages at the 5' end of a nucleic acid chain. This enzyme is localized to the mitochondria and may play a role in programmed cell death. Alternatively spliced transcript variants have been described. A pseudogene exists on chromosome 18. [provided by RefSeq, Feb 2009]
Applications:	ELISA, WB, IHC
Name of antibody:	EXOG
Immunogen:	Fusion protein of human EXOG
Full name:	exo/endonuclease G

Synonyms:	ENGL; ENGLA; ENGLB; ENGL-a; ENGL-b; ENDOGL1; ENDOGL2
SwissProt:	Q9Y2C4
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human colorectal cancer
IHC Recommend dilution:	150-300
WB Predicted band size:	41 kDa
WB Positive control:	K562, LOVO, HT-29, HUVEC cell lysates
WB Recommended dilution:	500-2000



