

兔抗 FABP3 多克隆抗体

中文名称: 兔抗 FABP3 多克隆抗体

英文名称: Anti-FABP3 rabbit polyclonal antibody

别 名: MDGI; FABP11; H-FABP; M-FABP; O-FABP

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: FABP3

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: Unconjugate

技术规格

st three distinct types, namely the hepatic-, intestinal- a nd cardiac-type. They form 14-15 kDa proteins and are thought to participate in the uptake, intracellular metab olism and/or transport of long-chain fatty acids. They m ay also be responsible in the modulation of cell growth and proliferation. Fatty acid-binding protein 3 gene cont ains four exons and its function is to arrest growth of

The intracellular fatty acid-binding proteins (FABPs) belongs to a multigene family. FABPs are divided into at lea

mammary epithelial cells. This gene is a candidate tumo



	r suppressor gene for human breast cancer.
Applications:	ELISA, WB, IHC
Name of antibody:	FABP3
Immunogen:	Synthetic peptide of human FABP3
Full name:	fatty acid binding protein 3, muscle and heart
Synonyms:	MDGI; FABP11; H-FABP; M-FABP; O-FABP
SwissProt:	P05413
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human brain and human lung cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	15 kDa
WB Positive control:	Human heart tissue
WB Recommended dilution:	200-1000









