

兔抗 FCAR 多克隆抗体

中文名称：兔抗 FCAR 多克隆抗体

英文名称： Anti-FCAR rabbit polyclonal antibody

别名： CD89

相关类别： 一抗

抗原： FCAR

储存： 冷冻（-20℃）

宿主： Rabbit

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

This gene is a member of the immunoglobulin gene superfamily and encodes a receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsinized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. M

	ultiple alternatively spliced transcript variants encoding different isoforms have been described for this gene.
Applications:	ELISA, WB, IHC
Name of antibody:	FCAR
Immunogen:	Fusion protein of human FCAR
Full name:	Immunoglobulin alpha Fc receptor
Synonyms:	CD89
SwissProt:	P24071
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human liver cancer and Human prostate cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	32 kDa
WB Positive control:	293T cells
WB Recommended dilution:	500-2000



