

## 兔抗 FCGR2A 多克隆抗体

中文名称：兔抗 FCGR2A 多克隆抗体

英文名称：Anti-FCGR2A rabbit polyclonal antibody

别名：CD32; CD32A; CDw32; FCG2; FCGR2; FCGR2A1; FcGR; IGFR2; MGC23887; MGC30032; FCGR2A

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：FCGR2A

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

#### Background:

CD32 (also designated Fc gamma RII) is a low affinity receptor for the Fc fragment of aggregated IgG. CD32 is responsible for the clearance of immunocomplexes by macrophages and also plays an important role in the regulation of antibody production by B cells. IgG can noncooperatively bind either one or two highly glycosylated CD32 molecules, and this binding delivers a negative signal for B cells. CD32 exists as several isoforms that are produced by alternative splicing of three distinct genes, A, B, and C. These isoforms are designated FcγRIIA, FcγRIIB1, FcγRIIB3,

	and FcγRIIC. All isoforms are present on monocytes, placental trophoblasts and endothelial cells. In addition, the FcγRIIB forms are present on B lymphocytes, and the FcγRIIA and FcγRIIC forms are found on neutrophils.
<b>Applications:</b>	WB
<b>Name of antibody:</b>	FCGR2A
<b>Immunogen:</b>	Fusion protein of human FCGR2A
<b>Full name:</b>	Fc fragment of IgG, low affinity IIa, receptor (CD32)
<b>Synonyms :</b>	CD32; CD32A; CDw32; FCG2; FCGR2; FCGR2A1; FcGR; IGFR2; MGC23887; MGC30032; FCGR2A
<b>SwissProt:</b>	P12318
<b>WB Predicted band size:</b>	35 kDa
<b>WB Positive control:</b>	Jurkat cells
<b>WB Recommended dilution:</b>	500-2000

