

兔抗 FCRLA 多克隆抗体

中文名称：兔抗 FCRLA 多克隆抗体

英文名称： Anti-FCRLA rabbit polyclonal antibody

别名： Fc receptor like A; FCRL; FCRX; FREB; FCRL1; FCRLX; FCRLb; FCRLd; FCRLe; FCRLM1; FCRLc1; FCRLc2

相关类别： 一抗

储存： 冷冻（-20℃）

宿主： Rabbit

抗原： FCRLA

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

This gene encodes a protein similar to receptors for the Fc fragment of gamma immunoglobulin (IgG). These receptors, referred to as FCGRs, mediate the destruction of IgG-coated antigens and of cells induced by antibodies. This encoded protein is selectively expressed in B cells, and may be involved in their development. This protein may also be involved in the development of lymphomas. Multiple alternatively spliced transcript variants that encode different protein isoforms have been described for this gene.

Applications:	ELISA, WB, IHC
Name of antibody:	FCRLA
Immunogen:	Fusion protein of human FCRLA
Full name:	Fc receptor like A
Synonyms:	FCRL; FCRX; FREB; FCRL1; FCRLX; FCRLb; FCRLd; FCRLe ; FCRLM1; FCRLc1; FCRLc2
SwissProt:	Q7L513
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human colorectal cancer and Human prostate cancer
IHC Recommend dilution:	100-200
WB Predicted band size:	39 kDa
WB Positive control:	Ramos cell lysate
WB Recommended dilution:	500-2000



