

兔抗 FGR 多克隆抗体

中文名称：兔抗 FGR 多克隆抗体

英文名称：Anti-FGR rabbit polyclonal antibody

别名：FGR proto-oncogene, Src family tyrosine kinase; SRC2; c-fgr; c-src2; p55-Fgr; p58-Fgr; p55c-fgr; p58c-fgr

抗原：FGR

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene is a member of the Src family of protein tyrosine kinases (PTKs). The encoded protein contains N-terminal sites for myristylation and palmitoylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to plasma membrane ruffles, and functions as a negative regulator of cell migration and adhesion triggered by the beta-2 integrin signal transduction pathway. I

	infection with Epstein-Barr virus results in the overexpression of this gene. Multiple alternatively spliced variants, encoding the same protein, have been identified.
Applications:	ELISA, IHC
Name of antibody:	FGR
Immunogen:	Synthetic peptide of human FGR
Full name:	FGR proto-oncogene, Src family tyrosine kinase
Synonyms:	SRC2; c-fgr; c-src2; p55-Fgr; p58-Fgr; p55c-fgr; p58c-fgr
SwissProt:	P09769
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	30-150

