

## 兔抗 MAP3K8(Ab-400) 多克隆抗体

中文名称：兔抗 MAP3K8(Ab-400) 多克隆抗体

英文名称： Anti-MAP3K8(Ab-400) rabbit polyclonal antibody

别名： COT; EST; ESTF; TPL2; MEKK8; Tpl-2; c-COT

相关类别： 一抗

储存： 冷冻（-20℃） 避光

宿主： Rabbit

抗原： MAP3K8(Ab-400)

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

### 技术规格

<b>Background:</b>	This gene is an oncogene that encodes a member of the serine/threonine protein kinase family. The encoded protein localizes to the cytoplasm and can activate both the MAP kinase and JNK kinase pathways. This protein was shown to activate I $\kappa$ B kinases, and thus induce the nuclear production of NF- $\kappa$ B. This protein was also found to promote the production of TNF- $\alpha$ and IL-2 during T lymphocyte activation. This gene may also utilize a downstream in-frame translation start codon, and thus produce an isoform containing a shorter N-terminus.
<b>Applications:</b>	WB, IHC

<b>Name of antibody:</b>	MAP3K8(Ab-400)
<b>Immunogen:</b>	Synthesized non-phosphopeptide derived from human MAP3K8 around the phosphorylation site of serine 400 (C-Q-S(p)-L-D).
<b>Full name:</b>	mitogen-activated protein kinase kinase kinase 8
<b>Synonyms :</b>	COT; EST; ESTF; TPL2; MEKK8; Tpl-2; c-COT
<b>SwissProt:</b>	P41279
<b>IHC positive control:</b>	Human brain tissue
<b>IHC Recommend dilution:</b>	50-100
<b>WB Predicted band size:</b>	53 kDa
<b>WB Positive control:</b>	COLO cells treated with Insulin (0.01U/ml, 15mins) lysate
<b>WB Recommended dilution:</b>	500-3000



