

## AMACR 抗原(重组蛋白)

- 中文名称: AMACR 抗原(重组蛋白)
- 英文名称: AMACR Antigen (Recombinant Protein)
- 储存: 冷冻(-20℃)
- 相关类别: 抗原

概述

Fusion protein corresponding to a region derived from 15-175 amino acids of human AMACR

技术规格

| Full name:         | alpha-methylacyl-CoA racemase  |
|--------------------|--|
| Synonyms:          | RM; RACE; CBAS4; AMACRD  |
| Swissprot:         | Q9UHK6   |
| Gene Accession:    | BC009471   |
| Purity:            | >85%, as determined by Coomassie blue stained SDS-PAGE   |
| Expression system: | Escherichia coli   |
| Tags:              | His tag C-Terminus, GST tag N-Terminus   |
| Background:        | This gene encodes a racemase. The encoded enzyme interconverts p<br>ristanoyl-CoA and C27-bile acylCoAs between their (R)- and (S)-stere<br>oisomers. The conversion to the (S)-stereoisomers is necessary for d<br>egradation of these substrates by peroxisomal beta-oxidation. Encod<br>ed proteins from this locus localize to both mitochondria and peroxi<br>somes. Mutations in this gene may be associated with adult-onset s<br>ensorimotor neuropathy, pigmentary retinopathy, and adrenomyelon<br>europathy due to defects in bile acid synthesis. Alternatively spliced<br>transcript variants have been described. Read-through transcription a<br>lso exists between this gene and the upstream neighboring C1QTNF |



| 3 ( | C1q | and | tumor | necrosis | factor | related | protein | 3) g | gene. |  |
|-----|-----|-----|-------|----------|--------|---------|---------|------|-------|--|
|     |     |     |       |          |        |         |         |      |       |  |