

兔抗 MAPK14 多克隆抗体

- 中文名称: 兔抗 MAPK14 多克隆抗体
- 英文名称: Anti-MAPK14 rabbit polyclonal antibody

别 名: mitogen-activated protein kinase 14; RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA

- 相关类别: 一抗
- 储 存: 冷冻 (-20℃)
- 宿 主: Rabbit
- 抗 原: MAPK14
- 反应种属: Human, Mouse, Rat
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

| Background: | The protein encoded by this gene is a member of the M |
|-------------|---|
| | AP kinase family. MAP kinases act as an integration point |
| | for multiple biochemical signals, and are involved in a wi |
| | de variety of cellular processes such as proliferation, diffe |
| | rentiation, transcription regulation and development. This |
| | kinase is activated by various environmental stresses and |
| | proinflammatory cytokines. The activation requires its pho |
| | sphorylation by MAP kinase kinases (MKKs), or its autoph |
| | osphorylation triggered by the interaction of MAP3K7IP1/ |
| | TAB1 protein with this kinase. The substrates of this kinas |



| | e include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related tr anscription and cell cycle regulation, as well as in genoto xic stress response. Four alternatively spliced transcript va riants of this gene encoding distinct isoforms have been reported. |
|-----------------------------|---|
| Applications: | ELISA, IHC |
| Name of antibody: | MAPK14 |
| Immunogen: | Fusion protein of human MAPK14 |
| Full name: | mitogen-activated protein kinase 14 |
| Synonyms: | RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14 ; PRKM15; SAPK2A; p38ALPHA |
| SwissProt: | Q16539 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human thyroid cancer |
| IHC Recommend dilution: | 100-300 |

