

BAG3 抗原（重组蛋白）

中文名称：BAG3 抗原（重组蛋白）

英文名称：BAG3 Antigen (Recombinant Protein)

储存：冷冻（-20℃）

相关类别：抗原

概述

Fusion protein corresponding to a region derived from 376-575 amino acids of human BAG3

技术规格

Full name:	BCL2 associated athanogene 3
Synonyms:	BIS; MFM6; BAG-3; CAIR-1
Swissprot:	O95817
Gene Accession:	BC014656
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The protein encoded by this gene contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner.

mlbio 梅联生物
Good elisakit producers