

兔抗 MOB2 多克隆抗体

中文名称：兔抗 MOB2 多克隆抗体

英文名称：Anti-MOB2 rabbit polyclonal antibody

别名：MOB kinase activator 2; HCCA2

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：MOB2

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

MOB2 (Mps one binder kinase activator-like 2), also known as HCCA2 (Hepatocellular carcinoma-associated gene 2), is a 467 amino acid protein that belongs to the MOB1/phocein family. MOB2 is expressed in lung, spleen, brain, and fetal liver. It is highly expressed in hepatocellular carcinoma. MOB2 is localized in the perinuclear region of the cytoplasm in liver cancer tissues and colocalizes with MAD2L2 in the nucleus of Hela cells. MOB2 is characterized by two N-glycosylation sites, six N-myristoylation sites, two Src homology 3 (SH3), and several phosphorylation motifs which indicate that this protein may pla

	<p>play a role in an intracellular signal transduction cascade. MOB2 binds to and regulates the autophosphorylation of the related human serine/threonine kinase 38 (NDR1) and serine/threonine kinase 38L(NDR2). It has been shown that MOB2 plays a critical role in cell cycle regulation. Overexpression of the protein during the G0/G1 phase inhibits cell proliferation causing cell cycle arrest. Stimulates the autophosphorylation and kinase activity of STK38 and STK38L.</p>
Applications:	ELISA, WB, IHC
Name of antibody:	MOB2
Immunogen:	Synthetic peptide of human MOB2
Full name:	MOB kinase activator 2
Synonyms:	HCCA2
SwissProt:	Q70IA6
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer and Human thyroid cancer
IHC Recommend dilution:	20-100
WB Predicted band size:	27 kDa
WB Positive control:	Mouse liver tissue and Rat liver tissue lysates
WB Recommended dilution:	200-1000



