

兔抗 GADL1 多克隆抗体

中文名称：兔抗 GADL1 多克隆抗体

英文名称：Anti-GADL1 rabbit polyclonal antibody

别名：glutamate decarboxylase like 1; ADC; CSADC; HuADC; HuCSADC

抗原：GADL1

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|------------------------------------|---|
| Background: | May catalyze the decarboxylation of aspartate, cysteine sulfinic acid, and cysteic acid to beta-alanine, hypotaurine and taurine, respectively. Does not exhibit any decarboxylation activity toward glutamate. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | GADL1 |
| Immunogen: | Synthetic peptide of human GADL1 |
| Full name: | glutamate decarboxylase like 1 |
| Synonyms: | ADC; CSADC; HuADC; HuCSADC |
| SwissProt: | Q6ZQY3 |
| ELISA Recommended dilution: | 5000-10000 |

| | |
|---------------------------------|----------------------------|
| IHC positive control: | Human cervical cancer |
| IHC Recommend dilution: | 30-150 |
| WB Predicted band size: | 59 kDa |
| WB Positive control: | 293T and K562 cell lysates |
| WB Recommended dilution: | 200-1000 |



