

兔抗 GCDH 多克隆抗体

- 中文名称:兔抗 GCDH 多克隆抗体
- 英文名称: Anti-GCDH rabbit polyclonal antibody
- 别名: glutaryl-CoA dehydrogenase; GCD; ACAD5
- 相关类别: 一抗
- 储存: 冷冻(-20℃)
- 抗原: GCDH
- 宿 主: Rabbit
- 反应种属: Human, Mouse
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

	The protein encoded by this gene belongs to the acyl-Co A dehydrogenase family. It catalyzes the oxidative decarbo xylation of glutaryl-CoA to crotonyl-CoA and CO(2) in the
	degradative pathway of L-lysine, L-hydroxylysine, and L-try
Background:	ptophan metabolism. It uses electron transfer flavoprotein as its electron acceptor. The enzyme exists in the mitocho
buckground.	ndrial matrix as a homotetramer of 45-kD subunits. Mutati
	ons in this gene result in the metabolic disorder glutaric a
	ciduria type 1, which is also known as glutaric acidemia ty
	pe I. Alternative splicing of this gene results in multiple tr
	anscript variants. A related pseudogene has been identifie



	d on chromosome 12.
Applications:	ELISA, IHC
Name of antibody:	GCDH
Immunogen:	Fusion protein of human GCDH
Full name:	glutaryl-CoA dehydrogenase
Synonyms:	GCD; ACAD5
SwissProt:	Q92947
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human cervical cancer
IHC Recommend dilution:	30-150

