

兔抗 HSD17B2 多克隆抗体

中文名称：兔抗 HSD17B2 多克隆抗体

英文名称：Anti-HSD17B2 rabbit polyclonal antibody

别名：HSD17; SDR9C2; EDH17B2

相关类别：一抗

储存：冷冻 (-20℃)

抗原：HSD17B2

宿主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	17 β -HSD2 (17 β hydroxysteroid dehydrogenase type 2) belongs to the 17 β -HSD family of proteins that regulate the availability of steroids within a tissue. 17 β -HSD2 converts active steroids to their inactive form through its oxidative activity. It is a key player in the inactivation of Estradiol and testosterone. Due to the effects that 17 β -HSD2 has on the availability of estrogen, it has been extensively investigated for playing a possible role in breast tumor development, colon cancer development and the pathophysiology of endometriosis.
Applications:	ELISA, WB, IHC
Name of antibody:	HSD17B2
Immunogen:	Fusion protein of human HSD17B2

Full name:	hydroxysteroid (17-beta) dehydrogenase 2
Synonyms :	HSD17; SDR9C2; EDH17B2
SwissProt:	P37059
ELISA Recommended dilution:	2000-5000
IHC positive control :	Human esophagus cancer and human brain
IHC Recommend dilution:	50-200
WB Predicted band size:	43 kDa
WB Positive control :	Human fetal liver tissue
WB Recommended dilution:	500-2000



