

兔抗 PTPN6(Ab-564) 多克隆抗体

- 中文名称: 兔抗 PTPN6(Ab-564) 多克隆抗体
- 英文名称: Anti-PTPN6(Ab-564) rabbit polyclonal antibody
- 别名: MSK1; RLPK; MSPK1
- 相关类别: 一抗
- 储存: 冷冻(-20℃)避光
- 宿 主: Rabbit
- 抗原: PTPN6(Ab-564)
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: Unconjugate

技术规格

Background:	Serine/threonine-protein kinase that is required f or the mitogen or stress-induced phosphorylatio n of the transcription factors CREB1 and ATF1 a nd for the regulation of the transcription factors RELA, STAT3 and ETV1/ER81, and that contribute s to gene activation by histone phosphorylation and functions in the regulation of inflammatory genes. Phosphorylates CREB1 and ATF1 in respo nse to mitogenic or stress stimuli such as UV-C irradiation, epidermal growth factor (EGF) and an
-------------	---



	isomycin. Plays an essential role in the control o
	f RELA transcriptional activity in response to TNF
	and upon glucocorticoid, associates in the cytopl
	asm with the glucocorticoid receptor NR3C1 and
	contributes to RELA inhibition and repression of
	inflammatory gene expression. In skeletal myobla
	sts is required for phosphorylation of RELA at 'S
	er-276' during oxidative stress. In erythropoietin-
	stimulated cells, is necessary for the 'Ser-727' ph
	osphorylation of STAT3 and regulation of its tra
	nscriptional potential. Phosphorylates ETV1/ER81
	at 'Ser-191' and 'Ser-216', and thereby regulates
	its ability to stimulate transcription, which may b
	e important during development and breast tum
	or formation. Directly represses transcription via
	phosphorylation of 'Ser-1' of histone H2A. Phosp
	horylates 'Ser-10' of histone H3 in response to
	mitogenics, stress stimuli and EGF, which results
	in the transcriptional activation of several immed
	iate early genes, including proto-oncogenes c-fo
	s/FOS and c-jun/JUN. May also phosphorylate 'S
	er-28' of histone H3. Mediates the mitogen- and
	stress-induced phosphorylation of high mobility
	group protein 1 (HMGN1/HMG14). In lipopolysac
	charide-stimulated primary macrophages, acts do
	wnstream of the Toll-like receptor TLR4 to limit
	the production of pro-inflammatory cytokines. Fu
	nctions probably by inducing transcription of the
	MAP kinase phosphatase DUSP1 and the anti-inf
	lammatory cytokine interleukin 10 (IL10), via CRE B1 and ATF1 transcription factors. Plays a role in
	neuronal cell death by mediating the downstrea
	m effects of excitotoxic injury.
Applications:	WB
Name of antibody:	PTPN6(Ab-564)
tanic of unibody.	Synthesized peptide derived from C-terminal of
Immunogen:	human SHP-1.
Full name:	ribosomal protein S6 kinase, 90kDa, polypeptide



	5
Synonyms :	MSK1; RLPK; MSPK1
SwissProt:	075582
WB Predicted band size:	90 kDa
WB Positive control:	Jurkat cells lysate
WB Recommended dilution:	500-3000