

兔抗 MUC1(Phospho-Tyr1229) 多克隆抗体

中文名称：兔抗 MUC1(Phospho-Tyr1229) 多克隆抗体

英文名称： Anti-MUC1(Phospho-Tyr1229) rabbit polyclonal antibody

别名： EMA; MCD; PEM; PUM; KL-6; MAM6; MCKD; PEMT; CD227; H23AG; MCKD1; MUC-1; ADMCKD; ADMCKD1; CA 15-3; MUC-1/X; MUC1/ZD; MUC-1/SEC

相关类别： 一抗

储存： 冷冻（-20℃） 避光

宿主： Rabbit

抗原： MUC1(Phospho-Tyr1229)

反应种属： Human

标记物： Unconjugate

克隆类型： Unconjugate

技术规格

Background:

This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex.

Applications:	WB, IHC, IF
Name of antibody:	MUC1(Phospho-Tyr1229)
Immunogen:	Peptide sequence around phosphorylation site of tyrosine 1229 S-P-Y(p)-E-K derived from Human CD227/MUC1.
Full name:	mucin 1, cell surface associated
Synonyms :	EMA; MCD; PEM; PUM; KL-6; MAM6; MCKD; PEMT; CD227; H23AG; MCKD1; MUC-1; ADMCKD; ADMCKD1; CA 15-3; MUC-1/X; MUC1/ZD; MUC-1/SEC
SwissProt:	P15941
IHC positive control:	Human breast carcinoma tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	122 kDa
WB Positive control:	HepG2 cells lysates treated with PMA
WB Recommended dilution:	500-1000
IF positive control:	HepG2 cells
IF Recommend dilution:	100-200



