

## 兔抗 ICAM1 多克隆抗体

- 中文名称：兔抗 ICAM1 多克隆抗体
- 英文名称：Anti-ICAM1 rabbit polyclonal antibody
- 别名：BB2; CD54; P3.58
- 相关类别：一抗
- 抗原：ICAM1
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 反应种属：Human
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

### 技术规格

<b>Background:</b>	This gene encodes a cell surface glycoprotein which is typically expressed on endothelial cells and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also exploited by Rhinovirus as a receptor. ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through ARHGEF26/SGEF and RHO G activation. In case of rhinovirus infection acts as a cellular receptor for the virus.
<b>Applications:</b>	ELISA, WB, IHC

<b>Name of antibody:</b>	ICAM1
<b>Immunogen:</b>	Synthetic peptide of human ICAM1
<b>Full name:</b>	intercellular adhesion molecule 1
<b>Synonyms :</b>	BB2; CD54; P3.58
<b>SwissProt:</b>	P05362
<b>ELISA Recommended dilution:</b>	2000-5000
<b>IHC positive control:</b>	Human lung cancer and Human breast cancer
<b>IHC Recommend dilution:</b>	50-200
<b>WB Predicted band size:</b>	58 kDa
<b>WB Positive control:</b>	Huvec, hela and raji cells
<b>WB Recommended dilution :</b>	500-2000



