

兔抗 ICAM3 多克隆抗体

- 中文名称：兔抗 ICAM3 多克隆抗体
- 英文名称：Anti-ICAM3 rabbit polyclonal antibody
- 别名：CD50; CDW50; ICAM-R
- 相关类别：一抗
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 抗原：ICAM3
- 反应种属：Human
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:	The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM) family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9 immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This protein is constitutively and abundantly expressed by all leucocytes and may be the most important ligand for LFA-1 in the initiation of the immune response. Its function is not only as an adhesion molecule, but also as a potent signaling molecule.
Applications:	ELISA, IHC
Name of antibody:	ICAM3

Immunogen:	Synthetic peptide of human ICAM3
Full name:	intercellular adhesion molecule 3
Synonyms:	CD50; CDW50; ICAM-R
SwissProt:	P32942
IHC positive control:	Human gastric cancer and human brain
IHC Recommend dilution:	25-100

